



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

Analisi dei femminicidi dalla casistica autoptica dell'Istituto di Medicina Legale di Bologna dal 1950 al 2019

Prof.ssa Susi Pelotti

Dipartimento di Medicina Legale – Università di Bologna



Contents lists available at [ScienceDirect](#)

Forensic Science International

journal homepage: www.elsevier.com/locate/forsciint



A 70-year study of femicides at the Forensic Medicine department, University of Bologna (Italy)

Arianna Giorgetti*, Paolo Fais, Elena Giovannini, Chiara Palazzo, Ilenia Filipuzzi,
Guido Pelletti, Susi Pelotti

Department of Medical and Surgical Sciences, Unit of Legal Medicine, University of Bologna, Via Irnerio 49, 40126 Bologna, Italy



Fonti

- Referti autoptici presso Archivio della Medicina Legale dell'Università di Bologna
- Referti autoptici delle CTU
- Archivi giornalistici nazionali online

Sono stati esclusi

- Infanticidio
- Morte accidentale
- Suicidio




Il nostro studio

- Età, nazionalità, attività lavorativa, stato di gravidanza
- Genere e relazione fra donna e aggressore
- Movente e luogo
- Causa e mezzi di produzione della morte
- Distribuzione anatomica e numero delle lesioni mortali



Risultati

172 omicidi di donne
di cui 103 femminicidi




classe d'età vittima	V.A.	%	x100mila
0-13	4	2,0%	0,11
14-17	3	1,5%	0,27
18-24	14	7,1%	0,71
25-34	25	12,7%	0,77
35-44	32	16,2%	0,77
45-54	40	20,3%	0,82
55-64	20	10,2%	0,48
65+	58	29,4%	0,75
non rilevato	1	0,5%	
TOT	197	100,0%	0,64

Decade	N.	Tasso per 100,000 abitanti (%)	Tasso per 100,000 abitanti genere femminile (%)	Età Media (SD)	Età Mediana (IR)
1950-59	28	0.37	0.71	37.8 (15.1)	32.5 (25.5-49)
1960-69	19	0.23	0.44	40.8 (16.2)	42 (29-49.5)
1970-79	33	0.36	0.69	38.3 (16.2)	35 (24-49)
1980-89	28	0.30	0.58	45.3 (22.6)	40 (30-61)
1990-99	21	0.23	0.45	46.9 (17.2)	47 (33-61)
2000-09	20	0.22	0.42	39.8 (24.2)	40 (22-54)
2010-19	23	0.23	0.45	60.5 (22.7)	58.5 (39-83)




Risultati


Nazionalità della donna
raccolta in 112 dei 172
casi (65%)



Risultati

Attività lavorativa
della donna raccolta in
51 dei 172 casi (30%)





The gender of the perpetrator was reported in 67% of the total sample and consisted in males for 90.4% and females for 9.6% of cases, showing no difference across the decades of observation.




Risultati

The relationship between the victim and the perpetrator was reported in 116 cases (67%) on the total sample of female homicides. Among these, 56.9% included an IP, either current husband or boyfriend (37.1%), previous husband or boyfriend (15.5%), or lover (4.3%). Perpetrator were family members in 23.3% of cases. In 17.2% of cases the perpetrator was an acquaintance of the victim, and in 3 cases (2.6%) it was reported that the victim and the perpetrator had no relationship at all (Fig. 2).

Among femicides, 63.1% were committed by IP, 18.5% by family partner, 17.5% by acquaintances and 0.97% by an unknown perpetrator (see Table A for more details).

Femminicidio commesso da:	V.A.	%
<i>partner</i>	113	57,4%
<i>ex partner</i>	25	12,7%
<i>figlio</i>	18	9,1%
<i>altro parente</i>	12	6,1%
<i>cliente / spacciatore</i>	9	4,6%
<i>padre</i>	9	4,6%
<i>altro conoscente</i>	5	2,5%
<i>autore non identificato</i>	5	2,5%
<i>autore sconosciuto alla vittima</i>	1	0,5%
TOT	197	100,0%



Association between motives for killing, age of the victim and victim-perpetrator relationship in the femicide sample. Percentages are separately calculated on column factors, i.e. the age of the victim and the victim-perpetrator relationship. IP: intimate partner.

	Age of the victim				Victim-perpetrator relationship			
	0-25 years (%)	26-45 years (%)	46-65 years (%)	> 65 years (%)	IP (%)	Family member (%)	Acquaintance (%)	No relationship (%)
Passionate reasons	25	52.9	9.1	7.7	44	0	11.1	0
Mental/physical disorder of the perpetrator	0	0	9.1	23.1	0	36.4	0	0
Mental disorder of the victim	0	0	0	46.1	12	36.4	0	0
Familial problems	25	17.7	36.4	23.1	20	27.3	22.2	0
Criminal motives	0	0	18.1	0	8	0	11.1	0
Others	50	29.4	27.3	0	16	0	55.6	0
Total	100	100	100	100	100	100	100	100


Table 3

Association between age of the victims, classified in four categories, and type of victim-perpetrator relationship in the femicide sample. Percentages are calculated on the basis of the column factor, i.e. victim-perpetrator relationship. IP: intimate partner.


		Victim-perpetrator relationship			
		IP (%)	Family (%)	Acquaintance (%)	No relationship (%)
Age of the victim	0-25 years	13.6	16.7	23.5	100
	26-45 years	45.8	11.1	29.4	0
	46-65 years	22	27.8	47.1	0
	> 65 years	18.6	44.4	0	0
Total		100	100	100	100




Causa della morte



Distribuzione anatomica della lesione fatale



Weapons or methods for murder



Modalità di morte della donna

