

- Abou-Kassem D.E., El-Kholy M.S., Alagawany M., Laudadio V., Tufarelli V. (2019). Age and sex-related differences in performance, carcass traits, hemato-biochemical parameters, and meat quality in Japanese quails. *Poultry Science* 98:1684-1691.
- Alam M.s., Howlider A.R., Mondal A., Hossain K., Rubayet Bostami A.B.M. (2008). Pattern of egg production in Japanese quail reared on littered floor and in cage. *Bangladesh Research Publications Journal*, 1(3): 239-249.
- Alkan S., Karsli T., Karabag K., Galiç A. (2013). The effects of different slaughter ages and sex on carcass characteristics in Japanese Quails of different lines (*Coturnix japonica*). *Süleyman Demirel Üniversitesi Ziraat Fakültesi Dergisi*, 8(1):12-18.
- Bayomy H.M., Rozan M.A., Mohammed G.M. (2017). Nutritional composition of quail meatballs and quail pickled eggs. *Journal of Nutrition & Food Sciences*, 7:2, DOI: 10.4172/2155-9600.1000584.
- Cheng K.M., Bennett D.C., Mills A.D. (2010). The Japanese Quail. *The UFAW Handbook on the Care and Management of Laboratory and other research animals*, VIII edition. <https://doi.org/10.1002/9781444318777.ch42>.
- Cullere M, Woods M J, van Emmenes L, Pieterse E, Hooffman L C, Dalle Zotte A (2019). *Hermetia illucens* larvae reared on different substrates in broiler quails dietary: effect on physicochemical and sensory quality of the quail meat. *Animals*, 9(8):525. doi.org/10.3390/ani9080525.
- Dalle Zotte A, Singh Y, Michiels J, Cullere M (2018). Black soldier fly (*Hermetia illucens*) as dietary source for laying quails: live performance, and egg physic – chemical quality, sensory profile and storage stability. *Animal* 12:3, 640-647.
- Dos Santos T.C., Gates R.S., de Fátima Ferriera Tinôco I., Estrada M.M., Chizzotti M.L. (2020). Meat quality traits of European quails reared under different conditions of temperature and air velocity. *Poultry Science* 99: 848-856.
- Karadağoğlu Ö., Şahin T., Ölmez M., Ahsan U., Özsoy B., Önk K. (2019). Fatty acid composition of liver and breast meat of quails fed diet containing black cumin (*Nigella sativa* L.) and/or coriander (*Coriandrum sativum* L.) seeds as unsaturated fatty acid sources. *Livestock Science*, 223, 164-171.
- Priti M., Satish S. (2014). Quail Farming: An Introduction. *Int. J. of Life Sciences*, 2014, Vol. 2(2): 190-193.
- Rezaei M., Karimi Torshizi M.A., Wall H., Ivarsson E. (2018). Body growth, intestinal morphology and microflora of quail on diets supplemented with micronized wheat fibre. *British Poultry Science*, 59:4, 422-429.
- Shanaway M.M. (1994). Quail production system. A review. *Food and Agriculture Organization of the United Nations*, Rome 1994.
- Tavaniello S. (2013). Effect of cross-breed of meat and egg line on productive performance and meat quality in Japanese quail (*Coturnix Japonica*) from different generations. - international Ph.D in Welfare, Biotechnology and quality of animal production.
- Thomas K.S., Jagatheesan R, Reetha T.L., Rajendran D. (2016). Nutrient composition of Japanese quail eggs. *International journal of Science Environment and Tecnology*, 5:3, 1293-1295.